

Author Index

- Abrams, R.M., Hutchison, A.A. and Gerhardt, K.J., Effect of high-intensity sound on local cerebral glucose utilization in fetal sheep, 1
- Albrecht, D. and Heinemann, U., Low calcium-induced epileptiform activity in hippocampal slices from infant rats, 316
- Alderson, R.F., Hua, Z.-W. and Hersh, L.B., Nerve growth factor and phorbol esters increase the number of choline acetyltransferase-positive cells in two morphologically distinct classes of basal forebrain neurons in primary cultures, 229
- Alesci, R., Raffaelli, A. and Bagnoli, P., Developmental changes of serotonin and 5-hydroxyindoleacetic acid levels in specific regions of the pigeon central nervous system, 151
- Bagnoli, P., see Alesci, R., 151
- Beazley, L.D., see Coleman, L.-A., 273
- Beazley, L.D., see Coleman, L.-A., 293
- Besson, J.M., see Dusart, I., 215
- Black, I.B., see Friedman, W.J., 177
- Blake, K., see Forloni, G., 73
- Blottner, D. and Unsicker, K., Spatial and temporal patterns of neurotrophic activities in rat adrenal medulla and cortex, 243
- Coleman, L.-A. and Beazley, L.D., Expanded retinofugal projections to the dorsal lateral geniculate nucleus and superior colliculus after unilateral enucleation in the wallaby *Setonix brachyurus*, quokka, 273
- Coleman, L.-A. and Beazley, L.D., Retinal ganglion cell number is unchanged in the remaining eye following early unilateral eye removal in the wallaby *Setonix brachyurus*, quokka, 293
- Cotman, C.W., see Peterson, C., 187
- Coyle, J.T., see Forloni, G., 73
- Curtis, E.M., Stewart, M.G. and King, T.S., Quantitation of synaptic, neuronal and glial development in the intermediate and medial hyperstriatum ventrale (IMHV) of the chick *Galus domesticus*, pre- and post-hatch, 105
- Dreyfus, C.F., see Friedman, W.J., 177
- Dusart, I., Nothias, F., Roudier, F., Besson, J.M. and Peschanski, M., Vascularization of fetal cell suspension grafts in the excitotoxically lesioned adult rat thalamus, 215
- Ferriero, D.M. and Sagar, S.M., Development of neuropeptide Y-immunoreactive neurons in the rat retina, 19
- Forloni, G., Blake, K., Hohmann, C.H. and Coyle, J.T., The postnatal expression of acetylcholinesterase in somatostatin-positive cells of mouse hippocampus, 73
- Friedman, W.J., Dreyfus, C.F., McEwen, B.S. and Black, I.B., Developmental regulation of tyrosine hydroxylase in the mediobasal hypothalamus, 177
- Gainer, H., see Szaro, B.G., 87
- Gerhardt, K.J., see Abrams, R.M., 1
- Gil-Loyzaga, P. and Parés-Herbute, N., HPLC detection of dopamine and noradrenaline in the cochlea of adult and developing rats, 157
- Gómez-Pinilla, F., Guthrie, K.M., Leon, M. and Nieto-Sampedro, M., NGF receptor increase in the olfactory bulb of the rat after early odor deprivation, 161
- Guthrie, K.M., see Gómez-Pinilla, F., 161
- Hafidi, A. and Romand, R., First appearance of Type II neurons during ontogenesis in the spiral ganglion of the rat. An immunocytochemical study, 143
- Heinemann, U., see Albrecht, D., 316
- Hersh, L.B., see Alderson, R.F., 229
- Hohmann, C.H., see Forloni, G., 73
- Hua, Z.-W., see Alderson, R.F., 229
- Hutchison, A.A., see Abrams, R.M., 1
- Kachele, D.L., see Lasiter, P.S., 27
- Kakarala, M.H., see Rovainen, C.M., 197
- Kawai, S., Yonetani, M., Nakamura, H. and Okada, Y., Effects of deprivation of oxygen and glucose on the neural activity and the level of high energy phosphates in the hippocampal slices of immature and adult rat, 11
- King, J.S., see Walker, J.J., 35
- King, T.S., see Curtis, E.M., 105
- Lasiter, P.S. and Kachele, D.L., Postnatal development of protein P-38 ('synaptophysin') immunoreactivity in pontine and medullary gustatory zones of rat, 27
- Laxer, U., see Yanai, J., 255
- Lee, V.M.-Y., see Szaro, B.G., 87
- Leon, M., see Gómez-Pinilla, F., 161
- McEwen, B.S., see Friedman, W.J., 177
- Mirrafati, S.J., see Robertson, R.T., 309
- Moore, S.J. and Munger, B.L., The early ontogeny of the afferent nerves and papillary ridges in human digital glabrous skin, 119
- Munger, B.L., see Moore, S.J., 119
- Nakamura, H., see Kawai, S., 11
- Neal, J.H., see Peterson, C., 187
- Nieto-Sampedro, M., see Gómez-Pinilla, F., 161
- Nothias, F., see Dusart, I., 215
- Okada, Y., see Kawai, S., 11
- Parés-Herbute, N., see Gil-Loyzaga, P., 157
- Peschanski, M., see Dusart, I., 215
- Peterson, C., Neal, J.H. and Cotman, C.W., Development of N-methyl-D-aspartate excitotoxicity in cultured hippocampal neurons, 187
- Pick, C.G., see Yanai, J., 255
- Poon, H.K., see Robertson, R.T., 309
- Raffaelli, A., see Alesci, R., 151
- Robertson, R.T., Poon, H.K., Mirrafati, S.J. and Yu, J., Transient patterns of acetylcholinesterase activity in developing thalamus: a comparative study in rodents, 309

- Romand, R., see Hafidi, A., 143
Roudier, F., see Dusart, I., 215
Rovainen, C.M. and Kakarala, M.H., Angiogenesis on the optic tectum of albino *Xenopus laevis* tadpoles, 197
Sagar, S.M., see Ferriero, D.M., 19
Schnitzer, J., see Wulle, I., 59
Senba, E., Yanaihara, C., Yanaihara, N. and Tohyama, M., Proenkephalin opioid peptide product in the sensory ganglia of the rat: a developmental immunohistochemical study, 263
Stewart, M.G., see Curtis, E.M., 105
Szaro, B.G., Lee, V.M.-Y. and Gainer, H., Spatial and temporal expression of phosphorylated and non-phosphorylated forms of neurofilament proteins in the developing nervous system of *Xenopus laevis*, 87
Tohyama, M., see Senba, E., 263
Trombka, D., see Yanai, J., 255
Unsicker, K., see Blottner, D., 243
Walker, J.J. And King, J.S., Ontogenesis of enkephalinergic afferent systems in the opossum cerebellum, 35
Wimer, C.C. and Wimer, R.E., On the sources of strain and sex differences in granule cell number in the dentate area of house mice, 167
Wimer, R.E., see Wimer, C.C., 167
Wulle, I. and Schnitzer, J., Distribution and morphology of tyrosine hydroxylase-immunoreactive neurons in the developing mouse retina, 59
Yanai, J., Laxer, U., Pick, C.G. and Trombka, D., Dopaminergic denervation reverses behavioral deficits induced by prenatal exposure to phenobarbital, 255
Yanaihara, C., see Senba, E., 263
Yanaihara, N., see Senba, E., 263
Yonetani, M., see Kawai, S., 11
Yu, J., see Robertson, R.T., 309

the 1990s, the number of people in the world who are undernourished has increased from 600 million to 800 million (FAO 1996).

There are a number of reasons for this increase. First, the world population has increased from 5 billion in 1987 to 6 billion in 1997, and is projected to reach 8 billion by 2025 (FAO 1996). Second, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996). Third, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996).

There are a number of reasons for this increase. First, the world population has increased from 5 billion in 1987 to 6 billion in 1997, and is projected to reach 8 billion by 2025 (FAO 1996). Second, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996). Third, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996).

There are a number of reasons for this increase. First, the world population has increased from 5 billion in 1987 to 6 billion in 1997, and is projected to reach 8 billion by 2025 (FAO 1996). Second, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996). Third, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996).

There are a number of reasons for this increase. First, the world population has increased from 5 billion in 1987 to 6 billion in 1997, and is projected to reach 8 billion by 2025 (FAO 1996). Second, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996). Third, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996).

There are a number of reasons for this increase. First, the world population has increased from 5 billion in 1987 to 6 billion in 1997, and is projected to reach 8 billion by 2025 (FAO 1996). Second, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996). Third, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996).

There are a number of reasons for this increase. First, the world population has increased from 5 billion in 1987 to 6 billion in 1997, and is projected to reach 8 billion by 2025 (FAO 1996). Second, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996). Third, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1996 (FAO 1996).